
This copy is for your personal, non-commercial use only.

If you wish to distribute this article to others, you can order high-quality copies for your colleagues, clients, or customers by [clicking here](#).

Permission to republish or repurpose articles or portions of articles can be obtained by following the guidelines [here](#).

The following resources related to this article are available online at www.sciencemag.org (this information is current as of December 23, 2011):

Updated information and services, including high-resolution figures, can be found in the online version of this article at:

<http://www.sciencemag.org/content/332/6032/930.full.html>

Supporting Online Material can be found at:

<http://www.sciencemag.org/content/suppl/2011/04/20/science.1203357.DC1.html>

This article **cites 42 articles**, 12 of which can be accessed free:

<http://www.sciencemag.org/content/332/6032/930.full.html#ref-list-1>

This article has been **cited by 5 articles** hosted by HighWire Press; see:

<http://www.sciencemag.org/content/332/6032/930.full.html#related-urls>

This article appears in the following **subject collections**:

Genetics

<http://www.sciencemag.org/cgi/collection/genetics>

Comparative Functional Genomics of the Fission Yeasts

Nicholas Rhind,^{1¶} Zehua Chen,² Moran Yassour,^{3,4,5¶} Dawn A. Thompson,^{3¶} Brian J. Haas,^{2¶} Naomi Habib,^{5,6¶} Ilan Wapinski,^{3,7¶} Sushmita Roy,^{3,8¶} Michael F. Lin,⁸ David I. Heiman,² Sarah K. Young,² Kanji Furuya,⁹ Yabin Guo,¹⁰ Alison Pidoux,¹¹ Hwei Mei Chen,¹² Barbara Robbertse,^{13*} Jonathan M. Goldberg,² Keita Aoki,⁹ Elizabeth H. Bayne,^{11†} Aaron M. Berlin,² Christopher A. Desjardins,² Edward Dobbs,¹¹ Livio Dukaj,¹ Lin Fan,² Michael G. FitzGerald,² Courtney French,⁶ Sharvari Gujja,² Klavs Hansen,^{14‡} Dan Keifenheim,¹ Joshua Z. Levin,² Rebecca A. Mosher,^{15§} Carolin A. Müller,¹⁶ Jenna Pfiffner,² Margaret Priest,² Carsten Russ,² Agata Smialowska,^{17,18} Peter Swoboda,¹⁷ Sean M. Sykes,² Matthew Vaughn,¹⁴ Sonya Vengrova,¹⁹ Ryan Yoder,¹³ Qiandong Zeng,² Robin Allshire,¹¹ David Baulcombe,¹⁵ Bruce W. Birren,²⁰ William Brown,¹⁶ Karl Ekwall,^{17,18} Manolis Kellis,^{8,3} Janet Leatherwood,¹² Henry Levin,¹⁰ Hanah Margalit,⁶ Rob Martienssen,¹⁴ Conrad A. Nieduszynski,¹⁶ Joseph W. Spatafora,¹³ Nir Friedman,^{5,21} Jacob Z. Dalgaard,¹⁹ Peter Baumann,^{22,23,24} Hironori Niki,⁹ Aviv Regev,^{3,4,24¶} Chad Nusbaum^{2¶}

The fission yeast clade—comprising *Schizosaccharomyces pombe*, *S. octosporus*, *S. cryophilus*, and *S. japonicus*—occupies the basal branch of Ascomycete fungi and is an important model of eukaryote biology. A comparative annotation of these genomes identified a near extinction of transposons and the associated innovation of transposon-free centromeres. Expression analysis established that meiotic genes are subject to antisense transcription during vegetative growth, which suggests a mechanism for their tight regulation. In addition, trans-acting regulators control new genes within the context of expanded functional modules for meiosis and stress response. Differences in gene content and regulation also explain why, unlike the budding yeast of Saccharomycotina, fission yeasts cannot use ethanol as a primary carbon source. These analyses elucidate the genome structure and gene regulation of fission yeast and provide tools for investigation across the *Schizosaccharomyces* clade.

The fission yeast genus *Schizosaccharomyces* forms a broad and ancient clade within the Ascomycete fungi (Fig. 1A) with a distinct life history from other yeasts (*I*). Fission yeast grow preferentially as haploids, divide by medial fission rather than asymmetric budding, and have evolved a single-celled life-style independently from the budding yeasts (Saccharomycotina). Fission yeasts share important biological processes with metazoans, including chromosome structure and metabolism—relatively large chromosomes, large repetitive centromeres, low-complexity replication origins, heterochromatic histone methylation, chromodomain heterochromatin proteins, small interfering RNA (siRNA)-regulated heterochromatin, and TRF family telomere-binding proteins—G₂/M cell cycle control, cytokinesis, the mitochondrial translation code, the RNA interference (RNAi) pathway, the signalosome pathway, and spliceosome components. These features are absent or highly diverged in budding yeast. In general, core orthologous genes in fission yeast more closely resemble those of metazoans than do those of other Ascomycetes (2). Fission yeasts have also evolved innovations in carbon metabolism, including aerobic fermentation of glucose to ethanol (3). This convergent evolution with the budding yeast *Saccharomyces cerevisiae* offers insight into the evolution of complex phenotypes.

S. pombe is widely used as a model for basic biological processes in the cell and to study genes implicated in human disease. To better understand its evolution and natural history, we have compared the genomes and transcriptomes of *S. pombe*, *S. japonicus*, *S. octosporus*, and *S. cryophilus*, which constitute all known fission yeasts.

Genome sequence and phylogeny. We sequenced and assembled the genomes of *S. octosporus*, *S. cryophilus*, and *S. japonicus* using clone-based and clone-free whole-genome shotgun (WGS) approaches (table S1). Each genome is ~11.5 Mb in size. *S. octosporus* and *S. cryophilus* are 38% GC; *S. japonicus* is 44%. By comparison, the *S. pombe* genome is 12.5 Mb in size and 36% GC. We assembled the *S. octosporus* and *S. japonicus* scaffolds into three full-length chromosomes of quality similar to that of the finished *S. pombe* genome (Fig. 1B, figs. S1 and S2, and tables S2 and S3) and identified telomeric sequence using WGS data (4). Telomere repeats in *S. japonicus* (GTCTTA), *S. octosporus* (GGGTACTT), and *S. cryophilus* (GGGTACTT) matched a one-and-a-half repeat-unit sequence at the putative telomerase-RNA locus, similar to the configuration in *S. pombe* (GGTTAC) (5). Using these motifs, we extended the *S. japonicus* and *S. octosporus* chromosomes into subtelomeric and telomeric sequence (4).

We constructed a phylogeny of the Schizosaccharomycetes within Ascomycota (Fig. 1A and fig. S3) from 440 single-copy core orthologs, placing the monophyletic *Schizosaccharomyces* species as a basal sister group to the clade, including the filamentous fungi (Pezizomycotina) and budding yeast (Saccharomycotina). We found an average amino acid identity of 55% between all 1:1 orthologs when we compared *S. pombe* and *S. japonicus*, similar to that between humans and the cephalochordate amphioxus (table S4). For the most closely related species, *S. cryophilus* and *S. octosporus*, 1:1 orthologs share 85% iden-

¹Biochemistry and Molecular Pharmacology, University of Massachusetts Medical School, 364 Plantation Street, Worcester, MA 01605, USA. ²Broad Institute of Massachusetts Institute of Technology and Harvard, 320 Charles Street, Cambridge, MA 02141, USA. ³Broad Institute of Massachusetts Institute of Technology and Harvard, 7 Cambridge Center, Cambridge, MA 02142, USA. ⁴Department of Biology, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139, USA. ⁵School of Computer Science and Engineering, Hebrew University, Jerusalem 91904, Israel. ⁶Department of Microbiology and Molecular Genetics, Faculty of Medicine, Hebrew University, Jerusalem 91120, Israel. ⁷Department of Systems Biology, Harvard Medical School, 200 Longwood Avenue, Alpert 536, Boston, MA 02115, USA. ⁸Computer Science and Artificial Intelligence Laboratory, Massachusetts Institute of Technology, 32 Vassar Street 32-D510, Cambridge, MA 02139, USA. ⁹Microbial Genetics Laboratory, Genetic Strains Research Center, National Institute of Genetics, 1111 Yata, Mishima, Shizuoka 411-8540, Japan. ¹⁰Eunice Kennedy Shriver National Institute of Child Health and Human Development, National Institutes of Health, Bethesda, MD 20892, USA. ¹¹Wellcome Trust Centre for Cell Biology, Institute of Cell Biology, School of Biological Sciences, The University of Edinburgh, 6.34 Swann Building, Mayfield Road, Edinburgh EH9 3JR, UK. ¹²Department of Molecular Genetics and Microbiology, Life Science, Room 130, State University of New York, Stony Brook, NY 11794, USA. ¹³Department of Botany and Plant Pathology, Oregon State University, Corvallis, OR 97331, USA. ¹⁴Cold Spring Harbor Laboratory, 1 Bungtown Road, Cold Spring Harbor, NY 11724, USA. ¹⁵Department of Plant Sciences, University of Cambridge, Downing Street, Cambridge CB2 3EA, UK. ¹⁶Centre for Genetics and Genomics, University of Nottingham, Queen's Medical Centre, Nottingham NG7 2UH, UK. ¹⁷Center for Biosciences, Department of Biosciences and Nutrition, Karolinska Institute, 141 38 Huddinge, Sweden. ¹⁸Department of Life Sciences, Södertörns Högskola, 141 89 Huddinge, Sweden. ¹⁹Warwick Medical School, University of Warwick, Gibbet Hill Campus, Coventry CV4 7AL, UK. ²⁰Broad Institute of Massachusetts Institute of Technology and Harvard, 301 Binney Street, Cambridge, MA 02141, USA. ²¹Alexander Silberman Institute of Life Sciences, Hebrew University, Jerusalem 91904, Israel. ²²Stowers Institute for Medical Research, Kansas City, MO 64110, USA. ²³Department of Molecular and Integrative Physiology, University of Kansas Medical School, Kansas City, KS 66160, USA. ²⁴Howard Hughes Medical Institute.

*Present address: National Center for Biotechnology Information, National Library of Medicine, National Institutes of Health, Department of Health and Human Services, 45 Center Drive, Bethesda, MD, 20892, USA.

†Present address: Wellcome Trust Centre for Gene Regulation and Expression, College of Life Sciences, University of Dundee, Dundee DD1 5EH, Scotland, UK.

‡Present address: Evolva Biotech A/S, Bülowvej 25, 1870 Frederiksberg C, Denmark.

§Present address: The University of Arizona, The School of Plant Sciences, 303 Forbes Building, 1140 East South Campus Drive, Tucson, AZ 87421, USA.

¶These authors made equivalent contributions.

¶¶To whom correspondence should be addressed. E-mail: nick.rhind@umassmed.edu (N.R.); aregev@broad.mit.edu (A.R.); chad@broadinstitute.org (C.N.)

tity on average, similar to humans and dogs. The genetic diversity within *S. pombe* is low. Comparing the *S. pombe* 972 strain to WGS analysis of *S. pombe* NCYC132 and *S. pombe* var. *kambucha*, two phenotypically distinct strains, revealed less than 1% nucleotide difference between the three strains (fig. S4 and table S5).

Eradication of transposons and reorganization of centromere structure. Transposons and other repetitive sequences are thought to be crucial for centromeric function through the maintenance of heterochromatin (6). These sequences evolve rapidly, but the evolutionary relation among centromeres, transposons, and heterochromatin is unclear, in part because fungal centromeres have not generally been included in genome assemblies. The *S. japonicus* genome harbors 10 families of gypsy-type retrotransposons (4) (fig. S5 and table S6). Sequence divergence of their re-

verse transcriptases suggests that these transposon families predate the last common ancestor of the Ascomycetes. However, a dramatic loss of transposons occurred after the divergence of *S. japonicus*; *S. pombe* harbors two related retrotransposons, Tf1 and Tf2; *S. cryophilus* has a single related retrotransposon, Tcry1; *S. octosporus* contains no transposons, but contains sequences related to reverse transcriptase and integrase that may represent extinct transposons (fig. S5 and table S6).

The disappearance of transposons in the evolution of fission yeast species after *S. japonicus* correlates with the appearance of the *cbp1* gene family, which suggests a transition in the control of centromere function. In *S. pombe*, Cbp1 proteins bind centromeric repeats and are required for transposon silencing and genome stability (7, 8). Although described as orthologs of CENP-B, a human centromere-binding protein, Cbp1 pro-

teins apparently evolved independently within the *Schizosaccharomyces* lineage from a domesticated Pogo-like DNA transposase (9). The appearance of the *cbp1* gene family also correlates with the switch from RNAi-mediated transposon silencing in *S. japonicus* (see below) to a Cbp1-based mechanism in *S. pombe*, which suggests that this shift to Cbp1-based transposon control allowed the eradication of most transposons from the fission yeast genomes, possibly by promoting recombinational deletion between long terminal repeats (LTRs) (8). Furthermore, the *cbp1* family is evolving rapidly (fig. S6), which suggests that Cbp1-based transposon silencing is a *Schizosaccharomyces*-specific innovation that arose after the divergence of *S. japonicus*.

The loss of transposons was accompanied by a substantial reorganization of chromosome architecture that conserves centromere function, which suggests that the evolution of novel




Fig. 1. *Schizosaccharomyces* phylogeny and chromosome structure. **(A)** A maximum-likelihood phylogeny of 12 fungal species from 440 core orthologs (each occurring once in each of the genomes) from fly to yeast. A maximum-parsimony analysis produces the same topology. Both approaches have 100% bootstrap support for all nodes. **(B)** The chromosome structure of *S. pombe*, *S. octosporus*, and *S. japonicus*. The middle bar in each figure represents the chromosome and its centromere: red for chromosome 1, blue for chromosome 2, and yellow for chromosome 3. Depicted above and below each chromosome are the

chromosomes in the other two species to which the genes on the chromosome of interest map, using the same color scheme. Depicted above the *S. pombe* and *S. japonicus* chromosomes are the distributions of transposons and mapping of siRNAs. *S. cryophilus* is not included, because its genome has not been assembled to complete chromosomes. **(C)** The centromeric repeat structures of *S. pombe*, *S. octosporus*, and *S. japonicus*. Due to their repetitive nature, they are unlikely to represent the exact genomic structure. The *S. pombe* portion of the figure is adapted from (11).

centromere structures compensated for the loss of transposons. In *S. japonicus*, transposons cluster next to telomeres and centromeres, as in metazoans (Fig. 1, B and C). In the other Schizosaccharomycetes, the subtelomeres and pericentromeres are also repetitive, but lack transposons (Fig. 1C). However, like *S. japonicus*, the centromeric and subtelomeric repeats are confined to pericentromeric and subtelomeric regions, respectively, with one exception—a centromeric repeat involved in transcriptional silencing at the *S. pombe* mating-type locus (10). We confirmed that the centromeres are heterochromatic by histone H3 lysine 9 (H3K9) methylation mapping (fig. S7) and showed that the *S. japonicus* centromeres are functional by meiotic mapping (table S2).

Although centromeric repeats evolve rapidly, differing even between related strains (11), individual repeat sequences tend to be similar within strains (Fig. 1C). No similarity was observed between the centromeric repeats of *S. pombe*, *S. octosporus*, or *S. cryophilus*. However, both *S. pombe* and *S. octosporus* centromeres contain repeated elements, highly similar between chromosomes, that are arrayed in a larger inverted repeat structure around a unique core sequence (Fig. 1C), which suggests that they are homogenized by nonreciprocal recombination. This contrasts with a lack of symmetry in *S. japonicus* and implies that transposition occurs more rapidly than homogenization by recombination. Thus, the suppression of transposition likely led both to the degeneration of transposon sequences and to the evolution of symmetric centromeric repeats.

Despite the divergence of centromere sequence and of gene order on the chromosome arms, karyotype and pericentromeric gene order are conserved between *S. pombe* and *S. octosporus* (fig. S8). Thus, although gene conversion maintains the similarity of centromeric repeats between the different centromeres, cross-over recombination between centromeres is suppressed. We observed neither centromeric translocations nor neocentromere events within these lineages, even though centromeres can occur at novel locations in manipulated *S. pombe* strains (12). The retention of repetitive elements in the centromeres of *S. pombe*, *S. octosporus*, and *S. cryophilus*, even as they have lost their transposons, implies that centromeric repeats have an important function.

Because siRNAs are involved in both transposon silencing and centromere function (13), we investigated these roles in the *Schizosaccharomyces* lineage. In *S. pombe*, the centromeric repeats produce dicer-dependent siRNAs required for maintenance of centromeric structure, function, and transcriptional silencing via Argonaute-dependent heterochromatin formation (14). However, transposons are silenced in *S. pombe* by RNAi-independent mechanisms and do not produce abundant siRNAs (Fig. 1B and fig. S9) (7). To investigate whether centromere-directed siRNA production is conserved within

the transposon-rich centromeres of *S. japonicus*, we sequenced small RNAs from log-phase *S. japonicus* cultures (which have a modal size of 23 nucleotides) (4) and found that 94% map to transposons, both telomeric and centromeric (Fig. 1B and fig. S9). The fact that siRNAs map to transposons in *S. japonicus* but not in *S. pombe* suggests that either the fission yeast RNAi pathway targets repetitive sequences instead of mobile elements per se or that the pathway evolved away from an ancestral role in transposon control to a dedicated role in heterochromatin function.

Evolution of mating-type loci. The structure of the mating-type loci and the cis-acting elements that regulate mating-type switching is highly conserved across all four species (fig. S10). The expressed *mat1* locus can contain either the plus (P) or the minus (M) allele and switches between the two by epigenetically programmed gene conversion (15–17) from two heterochromatically silenced donor cassettes: *mat2-P* and *mat3-M* (figs. S10 and S11). cis-Acting regulatory sequences required for epigenetic imprinting and recombinational switching (18–20) are conserved (4) (fig. S11), as is the epigenetically programmed genomic mark associated with *mat1* (15).

In contrast, none of the cis-acting sequences involved in transcriptional repression of the silent cassettes in *S. pombe* are identifiable in the

other species, although the donor cassettes are enriched for H3K9me heterochromatin (fig. S7). In *S. japonicus*, the silent *mat* loci are found to directly abut the centromere of chromosome 3, which suggests that they may be silenced by a positional effect. In *S. octosporus* and *S. cryophilus*, the *mat* loci are distant from the centromeres, but each contains a conserved region of transposon remnants, which may be silencing triggers. They also contain inverted repeats, albeit shorter and less similar to each other than the inverted repeats that flank the *mat2/3* locus in *S. pombe* (21). Thus, their silencing strategies may share elements from both *S. pombe* and *S. japonicus*. These results suggest that the mechanisms of imprinting and switching have been conserved, but that the strategies for establishing heterochromatin are plastic.

Comparative annotation of transcriptomes.

We annotated the three genomes using standard methods and compared them with *S. pombe* (4). We then deep-sequenced poly(A)-enriched, strand-specific cDNA (22–24) (RNA-Seq) and constructed de novo transcript models (fig. S12) for log phase, glucose depletion, early stationary phase, and heat shock from *S. pombe*, *S. octosporus*, and *S. japonicus* and log phase, glucose depletion, and heat shock from *S. cryophilus*.

In *S. pombe*, we reconstructed 4277 out of 5064 previously annotated genes; of the remain-




Fig. 2. Polyadenylated transcription is predominantly confined to protein-coding genes. The *S. pombe* genome was divided into four different feature classes: protein-coding sequence, intron sequence, untranslated sequence (5' and 3' UTRs), and intergenic sequence (all nucleotides between UTRs of protein-coding genes). The frequency of RNA-Seq reads was calculated over sequential 20-base pair windows across these features; for coding sequence, the frequency of antisense reads was also calculated. Frequency was normalized to the maximum frequency within each feature class to compensate for the different class sizes.

Table 1. Conservation of gene content and structure. See Fig. 5 legend for abbreviations.

Organism and total	Orthologous groups				Introns		
	Same	Gain	Loss	Dup	Same	Gain	Loss
<i>S. pombe</i>	4218	321	83	23	2901	297	27
<i>S. octosporus</i>	4218	133	48	5	2901	25	8
<i>S. cryophilus</i>	4218	283	73	11	2901	75	4
Ancestor of <i>Soct</i> and <i>Scry</i>	4218	103	44	15	2901	396	0
Ancestor of <i>Spom</i> , <i>Soct</i> , and <i>Scry</i>	4218	339	159	29	2901	415	412
<i>S. japonicus</i>	4218	242	0	18	2901	708	214
Ancestor of <i>Schizosaccharomyces</i>		640	745				
Total		2061	1152	101		1916	665

ing 788 genes, 60% were covered over at least 90% of their length. Four hundred of our transcript models change coding exon structure of the gene, 95% of which maintained or improved conserved coding capacity (tables S7 and S8 and fig. S12) (25). In addition, we identified 253 untranslated region (UTR) introns. Last, we found 89 new protein-coding genes in *S. pombe*, 53 of which are conserved (table S7 and fig. S13). We found no evidence that intron-rich fission yeast genes engage in metazoan-like alternative splicing (26). We found evidence for 433 alternative splicing events in *S. pombe* in the form of intron retention and alternative splice-donor or splice-acceptor usage, but no evidence of exon skipping or alternative exons; we found similar levels of splice variants in the other species (table S9). However, because many of these variants disrupt the coding capacity (figs. S14 and S15) and only a minority of intron retentions (146 out of 393) are conserved between two or more species, we suspect that much of alternative splicing in fission yeast represents nonproductive splicing variants. It is interesting that, in some cases, the nonspliced variant may be the protein-coding isoform (figs. S14, C and D, and S15, and table S10).

Transcription primarily represents protein-coding transcripts. The majority of stable fission yeast transcripts originate from annotated protein-coding genes. Most of the *S. pombe* genome is transcribed (22), with 91% of nu-

cleotides covered by at least one RNA-Seq read. However, most transcription, as measured by steady-state poly(A)-enriched RNA levels, is associated with well-defined transcripts, most of which are protein coding. Specifically, 37% of intergenic nucleotides (between the UTRs of annotated protein-coding transcripts) are not detectably expressed, and 90% of transcribed intergenic nucleotides account for only 0.16% of the poly(A)-enriched transcript signal. Moreover, the median expression level of exonic sequence (99.1% of which are detectably expressed) is 305 times that of intergenic sequence (Fig. 2 and table S11), with intergenic transcription enriched within origins of DNA replication (fig. S16)—gene-free loci with nucleosome-free regions (27–29) that may provide permissive loci for ectopic transcriptional initiation (30).

Transcription of coding genes is heavily biased to the sense strand. Of the coding genes, 73% have <5% of their RNA-Seq reads on the antisense strand. Genes with >5% antisense reads are enriched for convergent transcripts with intergenic distances of <200 bp ($P < 10^{-8}$, hypergeometric test), but not with those of >200 bp ($P > 0.1$), which suggests that much antisense transcription is due to readthrough of 3'-termination sites (31) (fig. S17). Thus, stable transcripts in fission yeast genomes are primarily associated with known transcription units. We discuss notable exceptions below.

Conservation of gene content and structure.

Despite the evolutionary breadth of the fission yeast clade, as measured by amino acid divergence, their gene content and structure are remarkably conserved. Of ~5000 coding genes in fission yeast species, 4218 are 1:1:1:1 orthologs across the clade, with the remainder of the orthologous groups containing genes that have been duplicated or deleted since their last common ancestor (Table 1 and fig. S12). Protein kinases are even more conserved in gene content; 93% (102 out of 110) of *S. pombe* protein kinases are 1:1:1:1 orthologs (4). Moreover, of 3601 *S. pombe* introns in 2616 spliced 1:1:1:1 orthologs, 2901 (81%) are identical across the four species (table S13). Overall, the conservation of gene content, gene order, and gene structure within *Schizosaccharomyces* is higher than expected given the level of amino acid divergence. From amino acid divergence, we estimate that the fission yeast clade arose about 220 million years ago (fig. S3). However, the conservation of gene content is significantly higher than that within *Saccharomyces* or *Kluyveromyces*, both of which have much lower amino acid divergence (table S15), which suggests that fission yeast amino acid sequences are evolving anomalously quickly or that genome structures are unusually stable.

The majority of gene changes are due to the gain of species- and clade-specific genes (table S12 and S14) (4). We tested whether gene gain is due to rapid divergence of orthologous genes by




Fig. 3. Meiotic genes are subject to antisense transcription. (A) Examples of antisense transcription of meiotic genes. Above and below the chromosome coordinates are the coding sequence annotations on the top and bottom strand, respectively. Above and below these are the strand-specific RNA-Seq read densities on a 0 to 300 scale; signal above 300 is truncated to make the low-amplitude signal visible. **(B)** Enrichment of Gene Ontology (GO) annotations within the set of protein-coding genes with more antisense than sense transcription. All terms with a P value of <0.01 are included, except for high-level terms (i.e., biological process and molecular function).

<i>S. pombe</i> GO annotation	p value
Meiosis	1.33×10^{-10}
Meiotic chromosome segregation	1.53×10^{-8}
Meiotic recombination	4.38×10^{-6}
Nuclear chromosome	4.54×10^{-3}
Ascospore formation	5.60×10^{-3}

looking for colinearity in regions with species-specific genes, and we examined these regions for signs of sequence similarity. We found that 94 out of 317 *S. pombe*-specific genes are in the same position relative to neighboring genes as genes specific to other species (table S16). Of these, nine show > 15% identity to a cognate gene in another species, which suggests that they are rapidly diverged orthologs (4).

We also found 34 *S. pombe* candidates for horizontal gene transfer from bacteria, including two published examples (4, 32, 33) (table S17), and similar numbers in the other species. Of these, 16 appear to have occurred before the radiation of the clade, and 9 appear to be specific to *S. pombe*.

Evidence for intergenic and antisense noncoding transcripts. We identified 1097 putative transcript models in *S. pombe* supported by strand-specific RNA-Seq data, but containing no obvious coding capacity and having no correspondence to well-defined noncoding RNAs (ncRNAs) (22, 24, 34) (fig. S18 and tables S18 and S19). Of these potential ncRNAs, 449 are intergenic and 648 are antisense, overlapping a coding gene on the other strand by at least 30%. Of the ncRNAs, 213 overlap an annotated UTR on the same strand, which suggests that they may be alternative UTRs. Nevertheless, the data support 338 of the intergenic and 546 of the antisense ncRNAs as distinct transcripts (4).

Of the 338 distinct intergenic ncRNAs in *S. pombe*, 138 are conserved in location in at least one other species (table S41). Moreover, 26 of the intergenic ncRNAs are conserved in sequence, and of these, 9 are conserved in both location and sequence, which suggests that they represent potentially biologically important noncoding RNAs. The transcripts that are conserved in location but not in sequence may represent functional transcripts that have diverged beyond recognition. Of the antisense transcripts, 328 (51%) are conserved across two or more genomes (table S20), which suggests that they are biologically important (35).

Antisense regulation of meiotic transcription. Across fission yeast, the ~250 genes with greater




Fig. 4. Expression profiles cluster into similar patterns with conserved biological functions. **(A)** Expression clusters for each species. Gene expression profiles for each species were clustered (4). The size of each heat map is proportional to the number of genes in the cluster and the number of genes in each is indicated. Similar cluster sizes and patterns reflect similar expression patterns between the species. The heat shock transcription profile is similar to log-phase growth because the tran-

scriptional response on the 15-min time scale used here is limited to a relatively small number of genes. L, log phase; GD, glucose depletion; ES, early stationary phase; HS, heat shock. **(B)** A selection of enriched GO terms for each cluster. The color intensity is proportional to the negative logarithm of the hypergeometric *P* value enrichment on a continuous scale of 0 to 10. Complete GO term enrichments are shown in table S26.

antisense transcription than sense transcription (table S21) are significantly enriched for meiotic genes ($P = 10^{-10}$ for *S. pombe*, hypergeometric test) (Fig. 3, fig. S19, and tables S22 and S23), consistent with observations in *S. pombe* and *S. cerevisiae* (24, 35). Several antisense-transcribed genes have been proposed to be regulated by intron retention (36, 37); however, these studies did not use strand-specific approaches, which makes it impossible to distinguish unspliced sense transcripts from antisense transcripts. We find no evidence of alternative splicing of any of these genes.

Antisense transcription of meiotic genes does not uniformly decrease as cognate sense transcription increases during meiosis (fig. S20). This observation suggests that antisense transcription does not inhibit sense transcription, in contrast to the anticorrelation observed in *S. cerevisiae* (30, 35). Furthermore, meiotic genes are not enriched among genes with >5% anti-

sense transcription but <100% antisense transcription ($P = 0.47$, hypergeometric test), consistent with a stoichiometric mechanism of regulation in which antisense transcripts directly bind to and inhibit the stability or translation of sense transcripts.

Global conservation of expression programs within fission yeasts. To identify conserved modules of coexpressed genes, we examined expression patterns across the four conditions and between the four fission yeast with phylogenetic clustering (Fig. 4). We found that patterns of gene expression between species grown in similar conditions are generally conserved, with dominant patterns associated with growth (log phase and heat shock) and stress (glucose depletion and early stationary phase). Moreover, similar expression clusters are enriched for similar gene annotations across the species.

Fission yeast up-regulate genes involved in mitosis, including those involved in the kine-

tochrome, the spindle pole body, and the anaphase-promoting complex, in response to glucose depletion (table S24). In contrast, several classes of genes involved in growth are down-regulated (4). None of these genes are extensively regulated in glucose depletion in *S. cerevisiae* (38).

cis-Regulatory mechanisms are associated with novel and expanded functions. Promoter motifs with conserved regulatory function across Ascomycota show new functionality among the *Schizosaccharomyces*. For example, the motif bound by Rtg3 in *S. cerevisiae* is associated with amino acid metabolism genes across the phylum. In fission yeast, however, it is also enriched in genes responsive to various stress responses (Fig. 5A). Of the stress genes that have Rtg3 motifs in *S. pombe*, 36% are found only in the *Schizosaccharomyces* clade, and many are also associated with the Atf1 motif, a conserved regulator of the stress response (Fig. 5B). Rtg3 does not have a detectable ortholog in the *Schizosaccharomyces*




Fig. 5. Conserved regulatory motifs have acquired new functions and new target genes. **(A)** The enrichment of gene functional modules regulated by the Rtg3-binding motif in 23 Ascomycota. This motif is enriched upstream of amino acid metabolism genes in all Ascomycota. However, in fission yeast it is specifically enriched upstream of stress-response genes. *S. cerevisiae* (Scer), *S. paradoxus* (Spar), *S. mikatae* (Smik), *S. bayanus* (Sbay), *C. glabrata* (Cgla), *S. castellii* (Scas), *K. waltii* (Kwal), *A. gossypii* (Agos), *K. lactis* (Klac), *S. kluyveri* (Sklu), *D. hansenii* (Dhan), *C. guilliermondii* (Cgui), *C. lusitanae* (Clus), *C. albicans* (Calb), *C. tropicalis* (Ctro), *C. parapsilosis* (Cpar), *C. elongosporus* (Celo), *Y. lipolytica* (Ylip), *N. crassa* (Ncra), *A. nidulans* (Anid), *S. japonicus* (Sjap), *S. octosporus* (Soct), and *S. pombe* (Spom). **(B)** Enrichment of Rtg3- and Atf1-binding sites in the promoters of stress-response genes. Each row represents a gene. The strength of the strongest regulatory site upstream of the gene is indicated in the blue heat map. The expression

of the gene in glucose depletion (gd) and early-stationary phase(es) relative to log phase is indicated in the blue-yellow heat map. Genes specific to the fission yeast clade are indicated in orange. **(C)** Enrichment of FKH- and MBF-binding sites in front of antisense-transcribed genes. As in (B), but each row represents a gene with greater antisense than sense transcription. Genes associated with meiosis (44) are indicated in magenta. CERSR, core environmental stress response.

clade (39), but the motif recognized by Rtg3 in *S. cerevisiae* is clearly identifiable in fission yeast, which suggests that these regulatory motifs are more conserved than their binding proteins. We also found a similar acquisition of *Schizosaccharomyces*-specific genes by the FKH- and MBF-associated motifs, which regulate meiotic transcription in *S. pombe* (4, 40, 41). In particular, these two motifs were enriched in genes with antisense transcripts (Fig. 5C). Most of the FKH (a motif bound by Mei4 in *S. pombe*) target genes with antisense transcripts (80%, 47 genes) are meiotic genes, the majority of which are specific to the *Schizosaccharomyces* clade (Fig. 5C).

Gene content reflects glucose-dependent lifestyle. Fission yeast and budding yeast of the *Saccharomyces* clade independently evolved the ability to produce ethanol by aerobic fermentation (3, 42). In contrast to the convergent evolution of ethanol production, the utilization of ethanol has not converged; although budding yeast can efficiently catabolize ethanol, fission yeast cannot use ethanol as a primary carbon source. The evolution of aerobic fermentation in budding yeast involved changes in gene content, most notably following a whole-genome duplication (WGD) event and in regulatory mechanisms of glucose repression (3, 43).

Like budding yeast, fission yeast have duplicate copies of the pyruvate decarboxylase (*pdc*) gene, needed to funnel pyruvate to fermentation. They also have orthologs of several activators and repressors of respiratory genes, including Hap2/3/4/5 complex members, the Adr1, Tup, and Mig transcriptional regulators, and the Snf1–Sip1 and 2 kinase (3). However, there are substantial distinctions in gene content between fission yeast and the post-WGD budding yeast (fig. S21). We identified loss of the glyoxylate cycle, loss of the glycogen biosynthesis, fewer glycolytic paralogs, loss of the gluconeogenic enzyme phosphoenolpyruvate carboxykinase, lack of expanded *adh* genes, and lack of transcriptional regulators of glucose repression as differences that illuminate the distinct metabolic capacities of fission yeast (4). All of these adaptations are consistent with the inability of fission yeast to consume ethanol as a sole carbon source. The loss of conserved enzymes highlights how fission yeast came to depend solely on glucose.

In both fission yeast and budding yeast, as glucose is depleted, the expression of respiratory genes [oxidative phosphorylation enzymes or tricarboxylic acid (TCA) cycle] is induced. However, unlike *S. cerevisiae* (38), in fission yeast the expression of the genes encoding the pyruvate dehydrogenase complex and *adh1* is reduced, which prevents the efficient use of pyruvate for respiration. Instead, the expression of the *ald* genes is induced, which may provide an alternative mechanism for generating acetyl-coenzyme A in fission yeast.

Thus, the lack of efficient ethanol catabolism by fission yeast demonstrates that aerobic

fermentation did not evolve to create a consumable by-product. Instead, ethanol is a waste product, possibly produced because it is toxic to competing microorganisms. It is interesting that aerobic fermentation appears to have evolved as early as 200 million years ago in fission yeast (fig. S3), long before the WGD and subsequent evolution of aerobic fermentation in budding yeast.

Conclusions. Our comparative analysis of genome structure and expression in the fission yeast, especially the analysis of centromere structure and evolution, demonstrates how chromosomal features can be rearranged while retaining function and maintaining stable positions across taxa. We also provide insight into centromeric biology and elucidate conserved antisense transcription that may play a systematic role in meiotic gene regulation. Last, this study informs our understanding of the major evolutionary innovation of aerobic alcohol fermentation in microbial metabolism that arose in parallel in the fission yeast and budding yeast lineages. As these results demonstrate, comparative analyses improve the power of fission yeast as a model for eukaryotic biology.

References and Notes

1. S. L. Forsburg, *Trends Genet.* **15**, 340 (1999).
2. V. Wood *et al.*, *Nature* **415**, 871 (2002).
3. C. L. Flores, C. Rodríguez, T. Petit, C. Gancedo, *FEMS Microbiol. Rev.* **24**, 507 (2000).
4. Materials and methods are available as supporting material on Science Online.
5. J. Leonardi, J. A. Box, J. T. Bunch, P. Baumann, *Nat. Struct. Mol. Biol.* **15**, 26 (2008).
6. L. H. Wong, K. H. Choo, *Trends Genet.* **20**, 611 (2004).
7. H. P. Cam, K. Noma, H. Ebina, H. L. Levin, S. I. Grewal, *Nature* **451**, 431 (2008).
8. M. Zaratiegui *et al.*, *Nature* **469**, 112 (2011).
9. C. Casola, D. Hucks, C. Feschotte, *Mol. Biol. Evol.* **25**, 29 (2008).
10. S. I. Grewal, A. J. Klar, *Genetics* **146**, 1221 (1997).
11. N. C. Steiner, K. M. Hahnenberger, L. Clarke, *Mol. Cell. Biol.* **13**, 4578 (1993).
12. K. Ishii *et al.*, *Science* **321**, 1088 (2008).
13. S. I. Grewal, *Curr. Opin. Genet. Dev.* **20**, 134 (2010).
14. T. A. Volpe *et al.*, *Science* **297**, 1833 (2002).
15. D. H. Beach, *Nature* **305**, 682 (1983).
16. R. Egel, *Curr. Genet.* **8**, 205 (1984).
17. S. Vengrova, J. Z. Dalgaard, *Genes Dev.* **18**, 794 (2004).
18. S. Sayrac, S. Vengrova, E. L. Godfrey, J. Z. Dalgaard, *PLoS Genet.* **7**, e1001328 (2011).
19. M. Kelly, J. Burke, M. Smith, A. Klar, D. Beach, *EMBO J.* **7**, 1537 (1988).
20. B. Arcangioli, A. J. Klar, *EMBO J.* **10**, 3025 (1991).
21. G. Singh, A. J. Klar, *Genetics* **162**, 591 (2002).
22. B. T. Wilhelm *et al.*, *Nature* **453**, 1239 (2008).
23. J. Z. Levin *et al.*, *Nat. Methods* **7**, 709 (2010).
24. T. Ni *et al.*, *PLoS ONE* **5**, e15271 (2010).
25. M. F. Lin, I. Jungreis, M. Kellis, *Nature Precedings*, published online 18 August 2010 (<http://hdl.handle.net/10101/npre.2010.4784.1>).
26. N. F. Käufer, J. Potashkin, *Nucleic Acids Res.* **28**, 3003 (2000).
27. M. Gómez, F. Antequera, *EMBO J.* **18**, 5683 (1999).

28. M. L. Eaton, K. Galani, S. Kang, S. P. Bell, D. M. MacAlpine, *Genes Dev.* **24**, 748 (2010).
29. A. B. Lantermann *et al.*, *Nat. Struct. Mol. Biol.* **17**, 251 (2010).
30. Z. Xu *et al.*, *Nature* **457**, 1033 (2009).
31. M. Zofall *et al.*, *Nature* **461**, 419 (2009).
32. T. Matsuzawa *et al.*, *Appl. Microbiol. Biotechnol.* **87**, 715 (2010).
33. T. Uo, T. Yoshimura, N. Tanaka, K. Takegawa, N. Esaki, *J. Bacteriol.* **183**, 2226 (2001).
34. N. Dutrow *et al.*, *Nat. Genet.* **40**, 977 (2008).
35. M. Yassour *et al.*, *Genome Biol.* **11**, R87 (2010).
36. A. Moldón *et al.*, *Nature* **455**, 997 (2008).
37. N. Averbek, S. Sunder, N. Sample, J. A. Wise, J. Leatherwood, *Mol. Cell* **18**, 491 (2005).
38. J. L. DeRisi, V. R. Iyer, P. O. Brown, *Science* **278**, 680 (1997).
39. I. Wapinski, A. Pfeffer, N. Friedman, A. Regev, *Nature* **449**, 54 (2007).
40. N. F. Lowndes, C. J. McInerney, A. L. Johnson, P. A. Fantes, L. H. Johnston, *Nature* **355**, 449 (1992).
41. H. Abe, C. Shimoda, *Genetics* **154**, 1497 (2000).
42. J. Piskur, E. Rozpedowska, S. Polakova, A. Merico, C. Compagno, *Trends Genet.* **22**, 183 (2006).
43. M. Kellis, N. Patterson, M. Endrizzi, B. Birren, E. S. Lander, *Nature* **423**, 241 (2003).
44. J. Mata, R. Lyne, G. Burns, J. Bähler, *Nat. Genet.* **32**, 143 (2002).

Acknowledgments: Assemblies and annotations are available at GenBank (*S. octosporus*, ABHY04000000; *S. cryophilus*, ACQJ02000000; *S. japonicus*, AATM02000000), the Broad Institute's *Schizosaccharomyces* Web site (www.broadinstitute.org/annotation/genome/schizosaccharomyces_group), which provides search and visualization tools and pomBase (www.pombase.org). The RNA-Seq and single-nucleotide polymorphism (SNP) data are at the National Center for Biological Information (NCBI), NIH, Sequence Read Archive (see table S42). The *S. japonicus* siRNA data sets are at NCBI GEO as GSE26902 and GSE27837. This work was supported by the National Human Genome Research Institute (NHGRI), NIH. C.N. and M.F.L. were supported by NHGRI; M.Y. was supported by a Clore Fellowship; I.W. is the Howard Hughes Medical Institute (HHMI) fellow of the Damon Runyon Cancer Research Foundation; S.R. was supported by NSF; R.M. was supported by NIH; K.H. was supported by Danish Research Council; C.A.N. and C.A.M. were supported by the Biotechnology and Biological Science Research Council; P.B. was supported by the Stowers Institute and HHMI; Y.G. and H.L. were supported by the National Institute of Child Health and Human Development, NIH; M.K. was supported by the NIH, an NSF CAREER award, and the Sloan Foundation; A.R. was supported by the Human Frontier Science Program, a Career Award at the Scientific Interface from the Burroughs Wellcome Fund, the Sloan Foundation, an NIH Director's PIONEER award and HHMI. We thank the Broad Institute Sequencing Platform; A. Fujiyama and A. Toyoda for generating DNA sequence; M. Lara and N. Stange-Thomann for developing molecular biology protocols; J. Robinson, M. Garber, and P. Müller for technical advice and support; A. Klar for providing *S. pombe* var. *kambucha* (SPK1820); L. Gaffney for assistance with the figures; K. Mar and J. Mwangi for administrative support; and C. Cuomo for comments on the manuscript.

Supporting Online Material

www.sciencemag.org/cgi/content/full/science.1203357/DC1
Materials and Methods
Figs. S1 to S25
Tables S1 to S42
References

26 January 2011; accepted 7 April 2011
Published online 21 April 2011;
10.1126/science.1203357